



A Higher Rate of Iron Deficiency in Obese Pregnant Sudanese Women

Wisal Abbas¹, Ishag Adam^{2*}, Duria A. Rayis², Nada G. Hassan², Mohamed F. Lutfi¹

¹Faculty of Medicine, Alneelain University, Khartoum, Sudan; ²Faculty of Medicine, University of Khartoum, P.O. Box 102, 11111, Khartoum, Sudan

Abstract

Citation: Abbas W, Adam I, Rayis DA, Hassan NG, Lutfi MF. A Higher Rate of Iron Deficiency in Obese Pregnant Sudanese Women. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2017.059>

Keywords: Body mass index; iron deficiency; obesity; pregnancy; Sudan.

***Correspondence:** Ishag Adam. Faculty of Medicine, University of Khartoum, P.O. Box 102, 11111, Khartoum, Sudan. Tel +249912168988. Fax +249183771211. E-mail: ishagadam@hotmail.com

Received: 08-Nov-2016; **Revised:** 10-Feb-2017; **Accepted:** 06-Mar-2017; **Online first:** 29-Apr-2017

Copyright: © 2017 Wisal Abbas, Ishag Adam, Duria A. Rayis, Nada G. Hassan, Mohamed F. Lutfi. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

Funding: This research did not receive any financial support.

Competing Interests: The authors have declared that no competing interests exist.

AIM: To assess the association between obesity and iron deficiency (ID).

MATERIAL AND METHODS: Pregnant women were recruited from Saad Abualila Hospital, Khartoum, Sudan, during January–April 2015. Medical history (age, parity, gestational age) was gathered using questionnaire. Weight and height were measured, and body mass index (BMI) was calculated. Women were sub-grouped based on BMI into underweight (< 18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25–29.9 kg/m²) and obese (≥ 30 kg/m²). Serum ferritin and red blood indices were measured in all studied women.

RESULTS: Two (0.5%), 126 (29.8%), 224 (53.0%) and 71 (16.8%) out of the 423 women were underweight, normal weight, overweight and obese, respectively. Anemia (Hb <11 g/dl), ID (ferritin <15µg/l) and iron deficiency anemia (IDA) were prevalent in 57.7%, 21.3% and 12.1%, respectively. Compared with the women with normal BMI, significantly fewer obese women were anemic [25 (35.2%) vs. 108 (85.7%), P < 0.001] and significantly higher number of obese women [25 (35.2) vs. 22 (17.5, P = 0.015) had iron deficiency. Linear regression analysis demonstrated a significant negative association between serum ferritin and BMI (– 0.010 µg/l, P= 0.006).

CONCLUSION: It is evident from the current findings that prevalence of anaemia and ID showed different trends about BMI of pregnant women.

Introduction

There is an increased rate of overweight and obesity among pregnant women [1]. The prevalence of overweight and obesity among Sudanese pregnant women at term are as high as 35.6% and 19.4%, respectively [2]. To our best of knowledge, Sudanese pregnant women have relatively higher body mass index (BMI) compared with other African countries especially in the Sub-Saharan region [2,3]. This creates a real challenge to obstetric practice because of the negative effects of obesity on the mother as well as the fetus [4]. The fetus of obese pregnant women is at higher risk of stillbirth, macrosomia, neural tube defects and other congenital anomalies [4–6]. Alternatively, gestational diabetes mellitus, preeclampsia, premature labour and caesarian delivery are more likely in obese compared with lean pregnant women [5,6].

Recent researches also revealed an intimate relation between high BMI and iron deficiency (ID) [7–9], adding more potential complications to obese pregnant women like iron deficiency anaemia (IDA) [9–14]. The negative impact of BMI on iron homeostasis was established in many studies targeting obese children [15,16] and adolescents [16,17]. In adults, studies exploring the link between high BMI and ID showed variable results, where significant associations were demonstrated in some reports [8,13] but not others [14,18]. Comparable studies among obese pregnant women are limited and mostly based on small sample size [19]. Likewise, studies assessing the impact of obesity on iron status during pregnancy did not consider the potential effect of the expected ID on erythropoiesis [10].

In this study, we assessed the association between high BMI and ID in pregnant Sudanese women. Also, the prevalence of IDA was evaluated in different BMI categories to judge the impact of ID on

erythropoiesis. The findings of the present study are expected to give more insight about iron homeostasis in obese pregnant women and to help health care providers in planning the proper intervention during antenatal visits.

Material and Methods

A cross-sectional study was conducted at Saad Abualila Maternity Hospital (Khartoum, Sudan) during the period January–April 2015. This is a tertiary hospital governed by the Faculty of Medicine, University of Khartoum. After signing an informed consent, eligible women were enrolled in the study. Inclusion criteria were: women with singleton pregnancy, at their early pregnancy (< 14 weeks of gestation), and willingness to participate in the study. Women with diabetes mellitus, thyroid disease, hypertension or any other chronic disease were excluded from the study. Obstetrics and medical history (age, parity, and gestational age) were gathered by a trained medical officer. Weight and height were measured, and BMI was calculated and expressed as weight in kilogrammes divided by the square of height in meters. The complete maternal hemogram with blood cell count was measured for all women, using an automated haematology analyser (Sysmex KX-21, Japan). Based on BMI, women were categorized into four groups: underweight (< 18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25–29.9 kg/m²) and obese (\geq 30 kg/m²). This classification follows the world health organisation, which addresses obesity during pregnancy as well as the general population [20]

Five ml of venous blood were taken from each participant, allowed to clot, centrifuged, and stored at –20°C until analysed. Serum ferritin was measured using radioimmunoassay gamma counter (Riostad, Germany) and kits provided by Beijing Isotope Nuclear Electronic Co., Beijing, China.

A total sample size of 430 participants was calculated using the previous incidence of obesity (19.4%) and incidence of IDA (29.4%) among pregnant women in Sudan [2], [21]. A formula was used to calculate the rate of iron deficiency in the obese and normal weight women, that would provide 80% power to detect a 5% difference at $\alpha = 0.05$, with an assumption that complete data might not be available for 10% of participants.

Statistics

SPSS for Windows (version 20.0) was used for data analyses. Studied variables were described with means (M), standard deviations (SD), median

and interquartile. Proportions of the studied groups were expressed in percentages (%). The difference of the continuous variables was compared between BMI groups using ANOVA when normally distributed (age, parity, gestational age, haemoglobin). Kruskal–Willis test was used for abnormally disturbed variables (serum ferritin). Linear regression analyses were performed, where BMI was the dependent variable and age, parity, gestational age, haemoglobin, and serum ferritin were the independent variables. $P < 0.05$ was considered statistically significant.

Ethics

The study received ethical clearance from the Research Board at the Department of Obstetrics and Gynaecology, Faculty of Medicine, University of Khartoum, Sudan.

Results

General characteristics of the 423 enrolled women are shown in Table 1. Around two fifths (176, 41.6%) of the women were primiparae. The mean (SD) of the age, parity and gestational age was 26.1 (6.2) years, 1.7 (2.2) and 10.7 (3.1) weeks, respectively.

Table 1: General characteristics of Sudanese women in the current study in their early pregnancy

| Variable | N = 423 |
|-----------------------------------|------------|
| <i>The mean (SD) of</i> | |
| Age, years | 26.8 (6.2) |
| Parity | 1.7 (2.2) |
| Gestational age, weeks | 10.7(3.1) |
| Body mass index kg/m ² | 26.6(3.7) |
| Haemoglobin, gm/dl | 10.5(0.8) |

Two (0.5%), 126 (29.8%), 224 (53.0%) and 71 (16.8%) out of the 423 women were underweight, normal weight, overweight, and obese, respectively.

Anaemia (Hb <11 g/dl) and iron deficiency (ferritin <15µg/l) were prevalent in 57.7% and 21.3% of the study sample, respectively. Of the total sample, 12.1% had iron-deficiency anaemia.

Although, there was no significant difference in the hemoglobin level and the other haematological parameters e.g. haematocrit (HCT), mean corpuscular volume (MCV), Mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) between women with normal BMI, underweight, overweight and obese, the median (interquartile) of serum ferritin level was significantly lower in obese compared with women with normal BMI, Table 2.

Compared with the women with normal BMI, significantly fewer obese women were anemic (25

[35.2%] vs. 108 [85.7%], $P < 0.001$) and significantly higher number of obese women (25 [35.2] vs. 22 [17.5], $P = 0.015$) had iron deficiency. There was no significant difference in the rate of iron deficiency anaemia (IDA) between women in the different BMI groups, Table 2.

Table 2: Comparison of socio-demographic, medical and obstetric characteristics between BMI groups among early-pregnancy Sudanese women

| Variables | Underweight N (2) | Normal N (126) | Overweight N (244) | Obese N (71) | P |
|---------------------------------|----------------------|-------------------|-----------------------|------------------|-------|
| <i>The mean (SD)</i> | | | | | |
| Age, years | 19.5 (6.3) | 24.9 (6.5) | 26.7 (6.1) | 26.4 (5.4) | 0.021 |
| Parity | 1.0 (1.4) | 1.7 (2.6) | 1.9 (2.1) | 1.6 (1.9) | 0.724 |
| Gestational age, weeks | 10.0 (2.8) | 10.6 (3.0) | 10.7 (3.2) | 11.0 (3.1) | 0.798 |
| Inter-pregnancy interval | - | 20.4 (12.3) | 11.4 (17.8) | 17.7 (26.2) | 0.007 |
| Hemoglobin, gm/dl | 11.1 (1.2) | 10.4 (0.8) | 10.5 (0.8) | 10.5 (0.8) | 0.628 |
| <i>Median and interquartile</i> | | | | | |
| Hematocrit,% | 31.0 (30.0) | 35.0 (32.1–37.5) | 35.0 (32.2–37.1) | 34.9 (32.2–37.1) | 0.503 |
| MCV, fl | 82.9 (78.7) | 83.3 (79.9–87.0) | 83.6 (79.5–86.9) | 82.3 (79.2–87.5) | 0.990 |
| MCH, pg | 28.0 (25.7) | 28.4 (26.6–30.1) | 28.3 (26.3–30.0) | 28.3 (26.2–30.0) | 0.858 |
| MCHC, g/dl | 33.7 (32.7) | 34.1 (33.0–35.0) | 34.0 (33.0–34.8) | 33.8 (33.0–34.9) | 0.721 |
| Ferritin, µg/l | 89.5 (36.0–89.5) | 39.0 (21.0–78.0) | 28.6 (15.9–53.6) | 24.0 (12.3–58.0) | 0.010 |
| <i>Number (%) of</i> | | | | | |
| Education ≤ secondary level | 2 (100) | 91 (72.2) | 153 (68.3) | 40 (56.3) | 0.076 |
| Anemia | 1 (50.0) | 108 (85.7) | 108 (48.2) | 25 (35.2) | < |
| Iron deficiency | 0 (0) | 22 (17.5) | 43 (19.2) | 25 (35.2) | 0.015 |
| Iron deficiency anemia | 0 (0) | 19 (15.1) | 24 (10.7) | 8 (11.3) | 0.619 |

Linear regression analysis demonstrated a significant negative association between serum ferritin and BMI ($-0.010 \mu\text{g/l}$, $P = 0.006$), Table 3.

Table 3: Linear regression analyses of the factors and body mass index in Khartoum, Sudan

| Variables | Coefficient | Standard error | P |
|-----------------|-------------|----------------|-------|
| Age | 0.099 | 0.033 | 0.003 |
| Parity | -0.167 | 0.090 | 0.063 |
| Gestational age | 0.068 | 0.057 | 0.230 |
| Hemoglobin | 0.098 | 0.217 | 0.650 |
| Serum ferritin | -0.010 | 0.004 | 0.006 |

Discussion

It is evident from the current findings that prevalence of anaemia and ID showed different trends in relation to BMI of pregnant women. According to the present results, anaemia is less common in overweight and obese pregnant women in spite of predominance of ID among these categories. Interestingly, the higher prevalence of ID among overweight and obese pregnant women does not induce a significant reduction of HCT, MCV, MCH and MCHC in these groups.

The association between high BMI and ID was documented in several previous reports [7, 9, 16, 22, 23], but not others [14, 18]. An inverse relationship between iron level and BMI was repeatedly demonstrated among children and adolescents [7, 16, 22, 23]; however, comparable studies in adults showed non-reproducible results [14, 18, 24].

In 321 children and adolescents, ID was noted in 38.8%, 12.1% and 4.4% of the studied obese, overweight and normal-weight subjects, respectively. In comparison, the percentages of IDA were 6.7%, 35% and 58.3% in the respective groups [7]. Similar findings were reproduced in at least three separate Iranian [16], Chinese [23] and Mexican [25] studies. Also, the cross-sectional data obtained from National Health and Nutrition Examination Survey III (1988-1994) confirms a twice risk of ID in high compared with low BMI children [22].

In adults, ID was documented among other micronutrient deficiencies in morbidly obese patients before bariatric surgery [8, 24]. A meta-analysis study evaluating iron status among overweight/obese subjects demonstrated an intimate relationship between ID and obesity, though haemoglobin level was higher in obese compared to normal weight adults [13]. The association analysis of the same study showed significant differences between studies investigating subjects under 18 years of age (Odds ratio = 1.78) and subjects 18 of age and older (Odds ratio = 0.92). This finding was further supported by Ogata *et al.*, who failed to demonstrate significant differences in iron concentration between obese and normal weight adult subjects [18]. In a separate study, decreased serum iron level was validated in overweight women, but not men [14].

High prevalence of ID among overweight/obese children and adolescents was attributed to insufficient iron intake, increased physical growth and blood volume in these age groups [7, 12, 25]. Alternatively, studies exploring iron homeostasis in obese women [11] and children [26] also demonstrated impaired iron absorption and consequently poor response to iron fortification. There are repeated evidence that impaired absorption of iron from the gastrointestinal tract (GIT) is secondary to increased expression of hepcidin [15, 17], which is a major inhibitor of iron absorption from GIT in humans [27]. Hepcidin was proved to correlate positively with leptin in iron deficient obese children, even after adjustment for possible confounders like BMI [15]. This suggests that enhanced hepcidin production, and consequently disturbed iron metabolism, is probably secondary to associated high leptin levels in obese subjects. Another hypothesis attributes high hepcidin levels to the mild inflammation associated with obesity [28].

The normal trend for hepcidin levels is to decrease progressively from the first to the third trimester [29]; however, the reverse was proved in obese pregnant women [10]. On the other hand, iron transfer across the placenta is enhanced during pregnancy even if maternal iron stores are depleted [30]. Studies on animal models [30] as well as humans [31] confirmed increased iron transfer across the placenta in ID states, probably secondary to enhanced expression of placental transferrin receptor 1 (pTFR1). This suggests that iron metabolism during

gestation is biased toward keeping an adequate iron supply to the fetus, even if mother iron stores were depleted. Based on the above narrative, the conjunction of increased hepcidin levels and pTFR1 expression is a likely explanation for ID in the obese pregnant women we studied. However, this hypothesis remains for further investigation by the researchers in the field.

Interestingly, the higher prevalence of ID among overweight and obese pregnant women we studied does not induce a significant reduction of HCT, MCV, MCH and MCHC in these groups. It was hypothesised that low grades of iron depletion might be asymptomatic [32] but progress gradually to iron-deficient erythropoiesis [33], and eventually IDA [33, 34]. Lack of significant reduction in red blood cells indices among obese pregnant we studied may be indicative of lower grades of ID in this group.

Noteworthy, hepcidin and pTFR1 levels were not estimated in the present report, which constituted potential study limitations. Evaluation of these parameters in future researches can help in a better understanding of the pathophysiology of ID among obese pregnant women.

The current findings obviously demonstrate paradoxical variations in the prevalence of anaemia and ID in obese pregnant women. Anaemia is infrequent in overweight and obese pregnant women in spite of their high risk of ID. The higher prevalence of ID among overweight/obese pregnant women we studied did not induce a significant reduction in red blood cells indices, suggesting lower grades of ID in these groups.

In conclusion, further studies that also evaluate hepcidin and pTFR1 levels are desirable for a better understanding of the pathophysiology of ID among obese pregnant women.

References

- Heslehurst N, Ells LJ, Simpson H, Batterham A, Wilkinson J, Summerbell CD. Trends in maternal obesity incidence rates, demographic predictors, and health inequalities in 36,821 women over a 15-year period. *BJOG*. 2007;114(2):187–94. <https://doi.org/10.1111/j.1471-0528.2006.01180.x> PMID:17305899
- Rayis DA, Abbaker AO, Salih Y, Diab TE, Adam I. Epidemiology of underweight and overweight-obesity among term pregnant Sudanese women. *BMC Res Notes*. 2010;3(1):327. <https://doi.org/10.1186/1756-0500-3-327> PMID:21134264 PMID:PMC3004927
- Villamor E, Msamanga G, Urassa W, Petraro P, Spiegelman D, Hunter DJ, et al. Trends in obesity, underweight, and wasting among women attending prenatal clinics in urban Tanzania, 1995-2004. *Am J Clin Nutr*. 2006;83(6):1387–94. PMID:16762951
- Thrift AP, Callaway LK. The effect of obesity on pregnancy outcomes among Australian Indigenous and non-Indigenous women. *Med J Aust*. 2014;201(10):592–5. <https://doi.org/10.5694/mja13.11170> PMID:25390266
- Leddy MA, Power ML, Schulkin J. The impact of maternal obesity on maternal and fetal health. *Rev Obstet Gynecol*. 2008;1(4):170–8. PMID:19173021 PMID:PMC2621047
- Cnattingius S, Villamor E, Johansson S, Edstedt Bonamy A-K, Persson M, Wikström A-K, et al. Maternal obesity and risk of preterm delivery. *JAMA*. 2013;309(22):2362–70. <https://doi.org/10.1001/jama.2013.6295> PMID:23757084
- Pinhas-Hamiel O, Newfield RS, Koren I, Agmon A, Lilos P, Phillip M. Greater prevalence of iron deficiency in overweight and obese children and adolescents. *Int J Obes Relat Metab Disord*. 2003;27(3):416–8. <https://doi.org/10.1038/sj.ijo.0802224> PMID:12629572
- Sánchez A, Rojas P, Basfi-Fer K, Carrasco F, Inostroza J, Codoceo J, et al. Micronutrient Deficiencies in Morbidly Obese Women Prior to Bariatric Surgery. *Obes Surg*. 2016;26(2):361–8. <https://doi.org/10.1007/s11695-015-1773-9> PMID:26108638
- Zhao L, Zhang X, Shen Y, Fang X, Wang Y, Wang F. Obesity and iron deficiency: a quantitative meta-analysis. *Obes Rev*. 2015;16(12):1081–93. <https://doi.org/10.1111/obr.12323> PMID:26395622
- García-Valdes L, Campoy C, Hayes H, Florido J, Rusanova I, Miranda MT, et al. The impact of maternal obesity on iron status, placental transferrin receptor expression and hepcidin expression in human pregnancy. *Int J Obes (Lond)*. 2015;39(4):571–8. <https://doi.org/10.1038/ijo.2015.3> PMID:25614087
- Mujica-Coopman MF, Brito A, López de Roma-a D, Pizarro F, Olivares M. Body mass index, iron absorption and iron status in childbearing age women. *J Trace Elem Med Biol*. 2015;30:215–9. <https://doi.org/10.1016/j.jtemb.2014.03.008> PMID:24813452
- Aigner E, Feldman A, Datz C. Obesity as an emerging risk factor for iron deficiency. *Nutrients*. 2014;6(9):3587–600. <https://doi.org/10.3390/nu6093587> PMID:25215659 PMID:PMC4179177
- Cheng HL, Bryant C, Cook R, O'Connor H, Rooney K, Steinbeck K. The relationship between obesity and hypoferraemia in adults: a systematic review. *Obes Rev*. 2012;13(2):150–61. <https://doi.org/10.1111/j.1467-789X.2011.00938.x> PMID:21981048
- Micozzi MS, Albanes D, Stevens RG. Relation of body size and composition to clinical biochemical and hematologic indices in US men and women. *Am J Clin Nutr*. 1989;50(6):1276–81. PMID:2596419
- del Giudice EM, Santoro N, Amato A, Brienza C, Calabrò P, Wiegierinck ET, et al. Hepcidin in obese children as a potential mediator of the association between obesity and iron deficiency. *J Clin Endocrinol Metab*. 2009;94(12):5102–7. <https://doi.org/10.1210/jc.2009-1361> PMID:19850683
- Moayeri H, Bidad K, Zadhoush S, Gholami N, Anari S. Increasing prevalence of iron deficiency in overweight and obese children and adolescents (Tehran Adolescent Obesity Study). *Eur J Pediatr*. 2006;165(11):813–4. <https://doi.org/10.1007/s00431-006-0178-0> PMID:16733667
- Aeberli I, Hurrell RF, Zimmermann MB. Overweight children have higher circulating hepcidin concentrations and lower iron status but have dietary iron intakes and bioavailability comparable with normal weight children. *Int J Obes (Lond)*. 2009;33(10):1111–7. <https://doi.org/10.1038/ijo.2009.146> PMID:19636315
- Ozata M, Mergen M, Oktenli C, Aydin A, Sanisoglu SY, Bolu E, et al. Increased oxidative stress and hypozincemia in male obesity. *Clin Biochem*. 2002;35(8):627–31. [https://doi.org/10.1016/S0009-9120\(02\)00363-6](https://doi.org/10.1016/S0009-9120(02)00363-6)
- Dao MC, Sen S, Iyer C, Klebenov D, Meydani SN. Obesity during pregnancy and fetal iron status: is Hepcidin the link? *J Perinatol*. 2013;33(3):177–81. <https://doi.org/10.1038/jp.2012.81> PMID:22722675 PMID:PMC3718280
- Ota E, Haruna M, Suzuki M, Anh DD, Tho LH, Tam NTT, et al. Maternal body mass index and gestational weight gain and their association with perinatal outcomes in Viet Nam. *Bull World Health*

- Organ. 2011;89(2):127–36. <https://doi.org/10.2471/BLT.10.077982> PMID:21346924 PMCID:PMC3040376
21. Mubarak N, Gasim GI, Khalafalla KE, Ali NI, Adam I. Helicobacter pylori, anemia, iron deficiency and thrombocytopenia among pregnant women at Khartoum, Sudan. *Trans R Soc Trop Med Hyg.* 2014;108(6):380–4. <https://doi.org/10.1093/trstmh/tru044> PMID:24686789
22. Nead KG, Halterman JS, Kaczorowski JM, Auinger P, Weitzman M. Overweight children and adolescents: a risk group for iron deficiency. *Pediatrics.* 2004;114(1):104–8. <https://doi.org/10.1542/peds.114.1.104> PMID:15231915
23. Shi Z, Lien N, Kumar BN, Dalen I, Holmboe-Ottesen G. The sociodemographic correlates of nutritional status of school adolescents in Jiangsu Province, China. *J Adolesc Health.* 2005;37(4):313–22. <https://doi.org/10.1016/j.jadohealth.2004.10.013> PMID:16182142
24. Lefebvre P, Letois F, Sultan A, Nocca D, Mura T, Galtier F. Nutrient deficiencies in patients with obesity considering bariatric surgery: a cross-sectional study. *Surg Obes Relat Dis.* 10(3):540–6. <https://doi.org/10.1016/j.soard.2013.10.003> PMID:24630922
25. Cepeda-Lopez AC, Osendarp SJ, Melse-Boonstra A, Aeberli I, Gonzalez-Salazar F, Feskens E, et al. Sharply higher rates of iron deficiency in obese Mexican women and children are predicted by obesity-related inflammation rather than by differences in dietary iron intake. *Am J Clin Nutr.* 2011 May;93(5):975–83. <https://doi.org/10.3945/ajcn.110.005439> PMID:21411619
26. Zimmermann MB, Zeder C, Muthayya S, Winichagoon P, Chaouki N, Aeberli I, et al. Adiposity in women and children from transition countries predicts decreased iron absorption, iron deficiency and a reduced response to iron fortification. *Int J Obes (Lond).* 2008 Jul;32(7):1098–104. <https://doi.org/10.1038/ijo.2008.43> PMID:18427564
27. Ganz T. Heparin, a key regulator of iron metabolism and mediator of anemia of inflammation. *Blood.* 2003 Aug;102(3):783–8. <https://doi.org/10.1182/blood-2003-03-0672> PMID:12663437
28. Andrews M, Soto N, Arredondo-Olguín M. Association between ferritin and hepcidin levels and inflammatory status in patients with type 2 diabetes mellitus and obesity. *Nutrition.* 2015 Jan;31(1):51–7. <https://doi.org/10.1016/j.nut.2014.04.019> PMID:25441587
29. van Santen S, Kroot JJC, Zijdeveld G, Wiegerinck ET, Spaanderman MEA, Swinkels DW. The iron regulatory hormone hepcidin is decreased in pregnancy: a prospective longitudinal study. *Clin Chem Lab Med.* 2013 Jul;51(7):1395–401. <https://doi.org/10.1515/cclm-2012-0576> PMID:23241678
30. Gambling L, Czopek A, Andersen HS, Holtrop G, Srai SKS, Krejpcio Z, et al. Fetal iron status regulates maternal iron metabolism during pregnancy in the rat. *Am J Physiol Regul Integr Comp Physiol.* 2009 Apr;296(4):R1063-70. <https://doi.org/10.1152/ajpregu.90793.2008> PMID:19176888
31. Li Y-Q, Yan H, Bai B. Change in iron transporter expression in human term placenta with different maternal iron status. *Eur J Obstet Gynecol Reprod Biol.* 2008 Sep;140(1):48–54. <https://doi.org/10.1016/j.ejogrb.2008.02.012> PMID:18586377
32. Cook JD, Baynes RD, Skikne BS. Iron deficiency and the measurement of iron status. *Nutr Res Rev.* 1992 Jan;5(1):198–202. <https://doi.org/10.1079/NRR19920014> PMID:19094320
33. Jaber L, Diamond G. The Lack of Association of Iron Deficiency With Anemia in First Graders. *J Pediatr Hematol Oncol.* 2015 Oct;37(7):524–8. <https://doi.org/10.1097/MPH.0000000000000423> PMID:26334429
34. Le Nguyen Bao K, Tran Thuy N, Nguyen Huu C, Khouw I, Deurenberg P. Anemia and Iron Deficiency in Vietnamese Children, 6 to 11 Years Old. *Asia Pac J Public Health.* 2016 Jul;28(5 Suppl):94S–102S. <https://doi.org/10.1177/1010539516641506> PMID:27052301