

# In Vitro Nematophagous Activity of Predatory Fungi on Infective Nematodes Larval Stage of Strongyloidea Family

Majid Zarrin<sup>1,2\*</sup>, Mahmoud Rahdar<sup>3</sup>, Farzad Poormohamadi<sup>2</sup>, Ali Rezaei-Matehkolaei<sup>2</sup>

<sup>1</sup>Health Research Institute, Infectious and Tropical Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; <sup>2</sup>Department of Medical Mycology, Medical School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; <sup>3</sup>Department of Medical Parasitology, Medical School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

## Abstract

**Citation:** Zarrin M, Rahdar M, Poormohamadi F, Rezaei-Matehkolaei A. In Vitro Nematophagous Activity of Predatory Fungi on Infective Nematodes Larval Stage of Strongyloidea Family. Open Access Maced J Med Sci. <https://doi.org/10.3889/oamjms.2017.064>

**Keywords:** nematophagous activity; fungi; Strongyloidea.

**\*Correspondence:** Majid Zarrin, Department of Medical Mycology, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: (+98-611) 3330074; Fax: (+98-611)3332036. E-mail: [mjzarrin@yahoo.co.uk](mailto:mjzarrin@yahoo.co.uk)

**Received:** 14-Feb-2017; **Revised:** 12-Apr-2017; **Accepted:** 18-Apr-2017; **Online first:** 11-Jun-2017

**Copyright:** © 2017 Majid Zarrin, Mahmoud Rahdar, Farzad Poormohamadi, Ali Rezaei-Matehkolaei. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

**Funding:** This work was based on an MSc thesis (Farzad Poormohamadi) which was supported by Health Research Institute, Infectious and Tropical Diseases Research Center, Jundishapur University of Medical Sciences, Ahvaz, Iran (Grant 92135).

**Competing Interests:** The authors have declared that no competing interests exist.

**AIM:** The main goal of the present research conducted to assess the in vitro activity of the nematophagous fungi *Duddingtonia flagrans*, *Fusarium solani*, *Verticillium chlamidosporium*, and *Trichoderma harzianum*.

**MATERIAL AND METHODS:** Four isolates of fungi including *D. flagrans*, *F. solani*, *V. chlamidosporium* and *T. harzianum* were used in this study. Horse faeces were used to provide the larvae stage of Strongyloidea family for the experiments.

**RESULTS:** *D. flagrans* was the most effective fungus to reduce the population of the larval nematodes. *D. flagrans* was able to kill 100% of larvae after 14 days of incubation. The significant effect was seen after 7 days of incubation, therefore, the live larvae was decreased to 9, 11, 19 and 25 for *D. flagrans*, *V. chlamidosporium*, *F. solani* and *T. harzianum*, respectively.

**CONCLUSION:** Our results illustrated that *D. flagrans* were most successful fungus for reducing the number of Strongyloidea family larva stage from horse faeces. Follow *D. flagrans*, the live larvae was significantly reduced for *V. chlamidosporium*, *F. solani* and *T. harzianum*, respectively.

## Introduction

Many nematodes are parasites on plants, animals and humans and able to cause significant economic losses and serious health problems. The nematodes' eggs deposited in the environment via the faeces of their host, and survive for long time periods and create an infection source for other animals and humans, and complete their biological cycle [1]. Gastrointestinal tract nematodes parasites of animals are the main issue that reduces the development of the livestock industry. Currently, the main technique for controlling nematodes population in livestock animals and field is chemotherapy against worm using benzimidazole carbamate drugs such as albendazole, mebendazole and cambendazole. However, in recent years there are many reports on resistance of nematodes against conventional anti parasites drugs and this method is expensive [2, 3]. Therefore, development of novel techniques is required to control

of nematodes for the reason that the conventional methods are based on the use of nematicidal agents that are related to main environmental and health problems. Some control programs have been developed to reduce the undesirable effects of gastrointestinal parasitism. A promising strategy for controlling of parasitic nematodes is using natural enemies against these parasites [4-7]. A group of largely soil-living fungi have nematophagous activity. The quantity of nematode-trapping fungal species present in a particular soil and the population densities of them can significantly be diverse. The highest densities are typically found in autumn and 30 cm the upper of soil [8].

Several fungal species have been investigated as potential agents for biological control. In predator fungi group, the genera *Duddingtonia*, *Arthrobotrys* and *Monacrosporium* show up for their efficient environmental control of parasite nematodes [9-11].

One of the most well-known fungi is *Duddingtonia flagrans* that are studied in several countries in the world [9, 11]. This fungus develops the three-dimensional nets and creates huge quantities of resistant spores. Chlamydospores increased the amount of fungus that passes throughout the gastrointestinal tract of sheep. Moreover, chlamydospores have the ability to germinate, make colonies in the faeces and destroy the fresh infective larvae and break off the life cycle of the parasite.

The main goal of the present research conducted to assess the in vitro activity of the nematophagous fungi *Duddingtonia flagrans*, *Fusarium solani*, *Verticillium chlamidosporium*, and *Trichoderma harzianum*.

## Materials and Methods

### Fungal strains

Four isolates of fungi included one from *D. flagrans*, one from *F. solani* (PTCC 5284), one from *V. chlamidosporium* (PTCC 5179) and one from *T. harzianum* (IBRC-M 30059) were incubated in Petri dishes containing potato dextrose agar at 25°C for 10 days. After growth of the isolates, a culture disks, 4 mm in diameter, were transferred to 9 cm diameter. Petri dishes containing 20 mL of 2% water agar medium and were incubated for 10 days.

### Preparation of Strongylidae Family Larval Stage

Horse faeces were used to provide the larvae stage for the experiments. The faeces were tested by light microscope to confirm that they were infected by egg parasite. The samples are incubated at room temperature for seven days to obtain third larvae stage. For collecting the larvae, Baermann apparatus method was used. Subsequently, 20 grammes of faeces from infected horses was added on funnel containing sterile cloth and water and was incubated at room temperature for 24 hrs. Finally, the larvae were collected from the narrow side of the funnel. The collected larvae were rinsed 5 times with PBS solution and centrifuged for 2 min at 500 rpm and then for preventing the bacteria and fungi growth, penicillin-streptomycin (100 IU/ml) (Sigma, Germany) and amphotericin B (fungizone/Bristol-Myers Squibb, Paris) were added and kept at 4°C until use.

### Co-culture of Fungi and Larvae

For removing the antibiotics, the larvae were washed with PBS solution for 10 times. One ml of

larvae suspension containing 100 third stage nematode larvae from *Strongylidae* family was separately added to 2% water-agar medium Petri-dishes containing the fungal cultures. On days 1, 2, 3, 7, 14 and 21, the live larvae were counted by light microscope (10x) and recorded. One hundred larvae were added to water agar Petri dish without fungi as a control group. All experiments were repeated for three times.

## Results

The results in Table 1 show that of all tested fungi, *D. flagrans* was the most effective fungus to reduce population of the larval nematodes.

**Table1: The nematophagus effects of studied fungi on third stage larvae population**

Fungi	Decrease of Larvae Population (%)					
	1 day	2 days	3 days	7 days	14 days	21 days
<i>D. flagrans</i>	58	45	30	9	0	0
<i>V. chlamidosporium</i>	62	56	38	11	4	0
<i>F. solani</i>	67	53	43	19	11	3
<i>T. harzianum</i>	69	58	50	25	17	8
Control	80	76	67	56	49	40

*D. flagrans* was able to kill 100% of larvae after 14 days of incubation. The number of the live nematodes larvae was decreased after challenging with *V. chlamidosporium*, *F. solani* and *T. harzianum* to 4, 11 and 17 respectively when incubated 14 days. *V. chlamidosporium* killed 100% of larvae after 21 days.




Figure 1: The trapped nematode larvae by *D. flagrans*

The significant effect was seen after 7 days of incubation, therefore, the live larvae was decreased to 9, 11, 19 and 25 for *D. flagrans*, *V. chlamidosporium*, *F. solani* and *T. harzianum* respectively. Figures 1 and 2 show the trapped larvae by 2 nematophagous fungi *D. flagrans* and *V. chlamidosporium*.



Figure 2: The trapped nematode larvae by *V. chlamidosporium*

most successful fungus for reducing the number of *Strongylidae* family larva stage from horse faeces. Follow *D. flagrans*; the live larvae was significantly reduced for *V. chlamidosporium*, *F. solani* and *T. harzianum* respectively.

However, *in vitro* studies on nematophagous property of fungi have limitations. They typically overrate the action of an agent by not allowing the larvae to escape. Lack of reproducing the interferences in soil and changes in the environment is another limitation of *in vitro* tests. Nevertheless, these methods have advantages, for example, a short assessment time and work between small physical spaces. Furthermore, the control of interaction between nematode and fungus is much easier. Further studies of the biological efficacy of nematophagous fungi in the field are required to obtain applicable strategy to control nematodes larvae contamination.

## Discussion

The most important purpose of biological control is to enhance the natural enemies of nematodes in environment subsequently as to decrease nematode density. Biological control of nematodes with fungi has been studied in several countries [12, 13].

The importance of the current work and similar studies is better understood when considered the free-living parasitic nematodes exist closely to our environment.

The *in vitro* efficacy of the genus *Monacrosporium* on Phyto nematodes, *Cooperia punctate* and *H. placei*, was demonstrated by Gomes et al. [14]. A study by Araújo et al. also confirmed the feasibility of using nematode-trapping fungus *Arthrobotrys robusta* in the biological control of parasite bovine gastrointestinal nematodes [15].

In our study, the use of *D. flagrans* for biological control of larva nematodes has great potential. The styles of traps have been investigated in detail in some predatory fungi. The adhesive network is the most extensive style used by nematode-trapping fungi.

Traps are the critical tools used by the nematode-trapping fungus to capture and kill nematodes [16]. Furthermore, traps are a significant marker for switching of nematode-trapping fungus from a saprophytic to a predacious lifestyle.

Our results illustrated that *D. flagrans* were

## Acknowledgments

This work was based on an M. Sc thesis (Farzad Poormohamadi) which was supported by Health Research Institute, Infectious and Tropical Diseases Research Center, Jundishapur University of Medical Sciences, Ahvaz, Iran (Grant 92135).

## References

- Vilela VL, Feitosa TF, Braga FR, Araújo JV, Santos Ad, Morais DF, Souto DV, Athayde AC. Coadministration of nematophagous fungi for biological control over gastrointestinal helminths in sheep in the semiarid region of northeastern Brazil. *Vet Parasitol.* 2016; 221:139-43. <https://doi.org/10.1016/j.vetpar.2016.03.027> PMID:27084486
- Pivoto FL, Machado FA, Anezi-Junior PA, Weber A, Cezar AS, Sangioni LA, Vogel FS. Improving liveweight gain of lambs infected by multidrug-resistant nematodes using a FECRT-based schedule of treatments. *Parasitol Res.* 2014; 113(6):2303-10. <https://doi.org/10.1007/s00436-014-3885-x> PMID:24744221
- Falzon LC, Menzies PI, Shakya KP, Jones-Bitton A, Vanleeuwen J, Avula J, Stewart H, Jansen JT, Taylor MA, Learmount J, Peregrine AS. Anthelmintic resistance in sheep flocks in Ontario, Canada. *Vet Parasitol.* 2013; 193(1-3):150-62. <https://doi.org/10.1016/j.vetpar.2012.11.014> PMID:23218224
- Xue JJ, Hou JG, Zhang YA, Wang CY, Wang Z, Yu JJ, Wang YB, Wang YZ, Wang QH, Sung CK. Optimization of storage condition for maintaining long-term viability of nematophagous fungus *Esteya vermicola* as biocontrol agent against pinewood nematode. *World J Microbiol Biotechnol.* 2014;30(11):2805-10. <https://doi.org/10.1007/s11274-014-1704-2> PMID:25070159
- Assis RC, Luns FD, de Araújo JV, Braga FR, Assis RL, Marcelino J, Freitas PC, Andrade MA. An isolate of the nematophagous fungus *Monacrosporium thaumasium* for the control of cattle trichostrongyles in south-eastern Brazil. *J Helminthol.* 2015;89(2):244-9. <https://doi.org/10.1017/S0022149X14000091> PMID:24622279

6. Braga FR, Carvalho RO, Silva AR, Araújo JV, Frassy LN, Lafisca A, Soares FE. Predatory capability of the nematophagous fungus *Arthrobotrys robusta* preserved in silica gel on infecting larvae of *Haemonchus contortus*. *Trop Anim Health Prod*. 2014; 46(3):571-4. <https://doi.org/10.1007/s11250-014-0544-2> PMID:24510197
7. Zarrin M, Rahdar M, Gholamian A. Biological Control of the Nematode Infective larvae of Trichostrongylidae Family with Filamentous Fungi. *Jundishapur J Microbiol*. 2015; 8(3): e17614. <https://doi.org/10.5812/ijm.17614> PMID:25893084 PMCID:PMC4397948
8. Persmark L, Banck A, Jansson H-B. Population dynamics of nematophagous fungi and nematodes in an arable soil – vertical and seasonal fluctuations. *Soil Biol Biochem*. 1996; 28:1005–1014. [https://doi.org/10.1016/0038-0717\(96\)00060-0](https://doi.org/10.1016/0038-0717(96)00060-0)
9. Buzatti A, de Paula Santos C, Fernandes MA, Yoshitani UY, Sprenger LK, dos Santos CD, Molento MB. *Duddingtonia flagrans* in the control of gastrointestinal nematodes of horses. *Exp Parasitol*. 2015;159:1-4. <https://doi.org/10.1016/j.exppara.2015.07.006> PMID:26208781
10. Silva ME, Braga FR, de Gives PM, Millán-Orozco J, Uriostegui MA, Marcelino LA, Soares FE, Araújo AL, Vargas TS, Aguiar AR, Senna T, Rodrigues MG, Froes FV, de Araújo JV. Fungal Antagonism Assessment of Predatory Species and Producers Metabolites and Their Effectiveness on *Haemonchus contortus* Infective Larvae. *Biomed Res Int*. 2015;2015: 241582. <https://doi.org/10.1155/2015/241582> PMID:26504791 PMCID:PMC4609344
11. Silva ME, Braga FR, Borges LA, Oliveira JM, Lima Wdos S, Guimarães MP, Araújo JV. Evaluation of the effectiveness of *Duddingtonia flagrans* and *Monacrosporium thaumasium* in the biological control of gastrointestinal nematodes in female bovines bred in the semiarid region. *Vet Res Commun*. 2014;38(2):101-6. <https://doi.org/10.1007/s11259-014-9590-5>
12. Degenkolb T, Vilcinskas A. Metabolites from nematophagous fungi and nematicidal natural products from fungi as alternatives for biological control. Part II: metabolites from nematophagous basidiomycetes and non-nematophagous fungi. *Appl Microbiol Biotechnol*. 2016; 100(9):3813-24. <https://doi.org/10.1007/s00253-015-7234-5> PMID:26728016 PMCID:PMC4824808
13. Xin B, Lin R, Shen B, Mao Z, Cheng X, Xie B. The complete mitochondrial genome of the nematophagous fungus *Lecanicillium saksenae*. *Mitochondrial DNA*. 2015; 28:1-2.
14. Gomes AP, Araújo JV, Ribeiro RC. Differential in vitro pathogenicity of predatory fungi of the genus *Monacrosporium* for phytonematodes, free-living nematodes and parasitic nematodes of cattle. *Braz J Med Biol Res*. 1999; 32(1):79-83. <https://doi.org/10.1590/S0100-879X1999000100012> PMID:10347773
15. Araújo JV, Gomes APS, Guimarães MP. Biological control of bovine gastrointestinal parasites in southeastern by the nematode-trapping fungus *Arthrobotrys robusta*. *Rev Bras Parasitol Vet*. 1998; 2: 117-122.
16. Hsueh YP, Mahanti P, Schroeder FC, Sternberg PW. Nematode-trapping fungi eavesdrop on nematode pheromones. *Curr Biol*. 2013; 23(1):83-6. <https://doi.org/10.1016/j.cub.2012.11.035> PMID:23246407 PMCID:PMC4047969